

Representability of $\underline{\mathrm{Aut}}_{\mathcal{F}}$ and $\underline{\mathrm{End}}_{\mathcal{F}}$

Niko Naumann

February 1, 2008

Abstract

Recently N. Nitsure showed that for a coherent sheaf $\mathcal{F}$ on a noetherian scheme the automorphism functor $\underline{\mathrm{Aut}}_{\mathcal{F}}$ is representable if and only if $\mathcal{F}$ is locally free. Here we remove the noetherian hypothesis and show that the same result holds for the endomorphism functor $\underline{\mathrm{End}}_{\mathcal{F}}$ even if one asks for representability by an algebraic space.

MSC2000: 14A25

1 Statement of results

1.1

Let X be a scheme and $\mathcal{F}$ a quasi-coherent $\mathcal{O}_X$ -module of finite presentation. We are interested in the representability of the following two functors on the category of X -schemes:

$$\begin{aligned}\underline{\mathrm{Aut}}_{\mathcal{F}}(X') &:= \mathrm{Aut}_{\mathcal{O}_{X'}}(f^*\mathcal{F}) \\ \underline{\mathrm{End}}_{\mathcal{F}}(X') &:= \mathrm{End}_{\mathcal{O}_{X'}}(f^*\mathcal{F})\end{aligned}$$

for $f : X' \rightarrow X$ an X -scheme.

The result is as follows:

Theorem 1 *Let X be a scheme and $\mathcal{F}$ a quasi-coherent $\mathcal{O}_X$ -module of finite presentation. Then the following are equivalent:*

- 1) $\mathcal{F}$ is locally free.
- 2) $\underline{\text{Aut}}_{\mathcal{F}}$ is representable by a scheme.
- 2') $\underline{\text{End}}_{\mathcal{F}}$ is representable by a scheme.

If X is locally noetherian, these conditions are also equivalent to the following:

- 3) $\underline{\text{Aut}}_{\mathcal{F}}$ is representable by an algebraic space.
- 3') $\underline{\text{End}}_{\mathcal{F}}$ is representable by an algebraic space.

1.2

The equivalence of 1) and 2) in theorem 1 in case X is noetherian is the main result of [N]. Our proof follows the ideas of *loc.cit.* closely. The main steps are contained in the following two lemmas:

Lemma 1 *Let A be a local ring and M a finitely presented A -module which is not free. Then there is a local homomorphism $A \rightarrow B$ such that*

$$M \otimes_A B \cong B^n \oplus (B/b)^m,$$

for some $0 \neq b \in B, b^2 = 0$ and $m \geq 1, n \geq 0$.

If A is noetherian, B can be chosen to be artin.

We observe that in the last statement of the lemma the noetherian hypothesis is indispensable: let $(B, \mathfrak{m})$ be a local ring such that there is $0 \neq b \in \bigcap_{n \geq 1} \mathfrak{m}^n$. Clearly $(b^2) \subsetneq (b)$, so after dividing out (b^2) one gets a ring B as in the lemma but for any local homomorphism $f : B \rightarrow C$ with C noetherian one clearly has $f(b) = 0$.

Lemma 2 *Let S be a scheme and $S_0 \subseteq S$ a closed subscheme defined by a nilpotent ideal sheaf. Assume X is a flat S -scheme and $f : X \rightarrow Y$ is an S -morphism such that $f \times \text{id}_{S_0}$ is an isomorphism. Then f is an isomorphism.*

1.3

In order to treat the representability of $\underline{\text{End}}_{\mathcal{F}}$ we will use the following observation:

Lemma 3 *Under the assumptions of 1.1 the obvious natural transformation of (set-valued) functors $\underline{\text{Aut}}_{\mathcal{F}} \rightarrow \underline{\text{End}}_{\mathcal{F}}$ is relatively representable by an open immersion.*

For completeness we also include a proof of the next lemma which is essentially lemma 5 of [N] and shows the relative representability of a “parabolic” sub-group functor:

Let X be a scheme and

$$(1) \quad 0 \longrightarrow \mathcal{F}' \longrightarrow \mathcal{F} \longrightarrow \mathcal{F}'' \longrightarrow 0$$

a short exact sequence of quasi-coherent $\mathcal{O}_X$ -modules with $\mathcal{F}'$ finitely presented and $\mathcal{F}''$ locally free. For any morphism $f : Y \rightarrow X$, the sequence $f^*((1))$ is exact because $\mathcal{F}''$ is in particular $\mathcal{O}_X$ -flat and it makes sense to consider

$$P(Y) := \{\alpha \in \text{Aut}_{\mathcal{O}_Y}(f^*\mathcal{F}) \mid \alpha(f^*\mathcal{F}') \subseteq f^*\mathcal{F}'\} \subseteq \underline{\text{Aut}}_{\mathcal{F}}(Y) .$$

Lemma 4 *In the above situation, the natural transformation $P \hookrightarrow \underline{\text{Aut}}_{\mathcal{F}}$ is relatively representable by a closed immersion.*

For basic facts about (relative) representability we refer to [BLR], 7.6.

2 Proofs

2.1

In this subsection we dispense with the easy implications of theorem 1, the assumptions and notations of which we now assume:

As $\underline{\text{Aut}}_{\mathcal{F}}$ and $\underline{\text{End}}_{\mathcal{F}}$ are clearly Zariski sheaves the problem of representing them is Zariski local on X , i.e. we can assume that X is affine and $\mathcal{F}$ corresponds to a free module of finite rank. In this case, representability of both $\underline{\text{Aut}}_{\mathcal{F}}$ and $\underline{\text{End}}_{\mathcal{F}}$ is obvious; we have proved the implications $1) \Rightarrow 2)$ and $1) \Rightarrow 2')$. Finally, the implications $2) \Rightarrow 3)$ and $2') \Rightarrow 3')$ are trivial.

2.2

Proof of lemma 1: Let $(A, \mathfrak{m})$ be a local ring and M a finitely presented A -module which is not free. We will find the required local homomorphism $A \rightarrow B$ as a suitable quotient of A :

Let

$$(2) \quad A^m \xrightarrow{\alpha} A^n \xrightarrow{\beta} M \rightarrow 0$$

be a minimal presentation of M , i.e. $n = \dim_k(M/\mathfrak{m}M)$ where $k := A/\mathfrak{m}$ is the residue field of A . Then M is free if and only if $\alpha = 0$: clearly $\alpha = 0$ is sufficient for freeness of M and conversely, if M is free, it is necessarily so of rank n , hence β is a surjective endomorphism of A^n which must be an isomorphism by a standard application of Nakayama's lemma, c.f. [M], thm. 2.4., hence $\alpha = 0$.

For any $J \subseteq \mathfrak{m}$, $(2) \otimes_A A/J$ is a minimal presentation of the A/J -module M/JM . If we denote by $I \subseteq A$ the ideal generated by the coefficients of any matrix representation of α and note that the minimality of (2) implies $I \subseteq \mathfrak{m}$ we find that M/JM is A/J -free if and only if $\alpha \otimes id_{A/J} = 0$ if and only if $I \subseteq J$. As M is not A -free we have $I \neq 0$ and as I is finitely generated we get $\mathfrak{m}I \subsetneq I$, again by Nakayama's lemma. By Zorn's lemma, using again that I is finitely generated, there is an ideal J with $\mathfrak{m}I \subseteq J \subsetneq I$ and which is maximal subject to these conditions (indeed, any ascending chain of such ideals admits its union as an upper bound because I is finitely generated).

We claim that $B := A/J$ is as required:

By the maximality of J the ideal $\bar{I} := I/J$ is non-zero principal: $\bar{I} = (b)$, $0 \neq b \in B$ and we necessarily have $b^2 = 0$: if not, we would have $b \in (b^2)$, i.e. $b = xb^2$ or $b(1 - xb) = 0$ for some $x \in B$. As $b \in \bar{\mathfrak{m}} := \mathfrak{m}/J$, the maximal ideal of B , $1 - xb$ was a unit of B , so we would have $b = 0$.

We now show that $M \otimes_A B$ has the desired structure: any coefficient α_{ij} of

a matrix representation of $\alpha \otimes \text{id}_B$ is of the form $\alpha_{ij} = bu_{ij}, u_{ij} \in B$. As by construction $\overline{\mathfrak{m}}b = 0$ we see that if $\alpha_{ij} \neq 0$, then $u_{ij} \in B^*$. We get a matrix equation $(\alpha_{ij}) = b(u_{ij})$ and (u_{ij}) can be chosen with $u_{ij} = 0$ or $u_{ij} \in B^*$, all i, j . Then the usual Gauß-algorithm can be applied to (u_{ij}) , showing that indeed $M \otimes_A B \cong B^n \oplus (B/b)^m$ for some $m, n \geq 0$. As, by construction, $M \otimes_A B$ is not B -free, we finally see that $m \geq 1$.

If A is noetherian we can start the construction of B by first dividing out a suitable high power of $\mathfrak{m}$: Indeed, if $M/\mathfrak{m}^n M$ was free for all $n \geq 1$ we would have $I \subseteq \bigcap_{n \geq 1} \mathfrak{m}^n = (0)$. Then the ring B we obtain in the above construction is noetherian local with $\overline{\mathfrak{m}}$ nilpotent, hence zero-dimensional, i.e. B is artin local.

Proof of lemma 2: We can assume that the ideal sheaf $\mathcal{I}$ of $S_0 \subseteq S$ satisfies $\mathcal{I}^2 = 0$. Our assertion is local on S, X and Y and thus reduces to the following:

Given a ring k and an ideal $I \subseteq k$ of square zero, if $f : A \rightarrow B$ is a morphism of k -algebras with B k -flat and such that $f \otimes_k \text{id}_{k/I}$ is an isomorphism, then f is an isomorphism:

1) f is surjective: any $b \in B$ can be written

$$b = f(a) + \sum_j \alpha_j b'_j, \text{ some } \alpha_j \in I, b'_j \in B, a \in A.$$

Applying this to the b'_j we get (for some $\alpha_{ij} \in I, b''_{ij} \in B, a_j \in A$):

$$b = f(a) + \sum_j \alpha_j (f(a_j) + \sum_{ij} \alpha_{ij} b''_{ij}) = f(a + \sum_j \alpha_j a_j).$$

2) f is injective: For $K := \ker(f)$ the k -flatness of B implies $K/IK = 0$ and the same argument as in 1) shows that $K = 0$.

Proof of 2) $\Rightarrow$ 1) in theorem 1: Under the notations of 1.1 we assume that $\underline{\text{Aut}}_{\mathcal{F}}$ is representable by a scheme and, by contradiction, that $\mathcal{F}$ is not locally free. Note that the assumption on representability is stable under base-change $Y \rightarrow X$. So, base-changing to a suitable local ring of X , we can assume $X = \text{Spec}(A)$ with A local and $\mathcal{F}$ corresponding to a finitely presented A -module M which is not free. According to lemma 1 we can assume $M \cong A^n \oplus (A/a)^m$ for some $0 \neq a \in A$ with $a^2 = 0$ and $m \geq 1$. Let $G \rightarrow S := \text{Spec}(A)$ be the group-scheme representing $\underline{\text{Aut}}_M$ and put

$S_0 := \operatorname{Spec}(A/a)$. The sub-functor $G' \hookrightarrow G$ of automorphisms preserving (base-changes of) the direct summand $(A/a)^m$ is represented by a closed sub-group scheme (still to be denoted G') according to lemma 4.

Let $P \subseteq \operatorname{Gl}_{n+m,S}$ denote the standard parabolic sub-group of automorphisms preserving the rank m direct summand. P is flat over S , as can be seen over $\operatorname{Spec}(\mathbb{Z})$. There is a morphism of S -groups $f : P \rightarrow G'$ which on points is given by sending $\begin{pmatrix} \alpha & 0 \\ \beta & \gamma \end{pmatrix}$ to $\begin{pmatrix} \alpha & 0 \\ \pi\beta & \bar{\gamma} \end{pmatrix}$, where $\alpha \in \operatorname{Aut}_A(A^n)$, $\gamma \in \operatorname{Aut}_A(A^m)$, $\beta : A^n \rightarrow A^m$ and $\bar{\gamma} \in \operatorname{Aut}_{A/a}((A/a)^m)$ denotes the reduction of γ and $\pi : A^m \rightarrow (A/a)^m$ is the natural map. This “point-wise” description of f is immediately checked to be functorial and a homomorphism and hence does indeed define a morphism of S -groups. Obviously, $f \times \operatorname{id}_{S_0}$ is an isomorphism, hence so is f by lemma 2. This is however a contradiction, because $f(S) : P(S) \rightarrow G'(S)$ is not injective, as $f(S)(\operatorname{id}_{A^n} \oplus (1-a)\operatorname{id}_{A^m}) = 1$ and $a \neq 0, m \geq 1$.

2.3

Proof of lemma 3: Given a scheme X , a quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$ of finite presentation and some $\varphi \in \operatorname{End}_{\mathcal{O}_X}(\mathcal{F})$ we have to show that there is an open sub-scheme $X_0 \subseteq X$ such for all $f : Y \rightarrow X$, $f^*(\varphi) \in \operatorname{Aut}_{\mathcal{O}_Y}(f^*\mathcal{F}) \subseteq \operatorname{End}_{\mathcal{O}_Y}(f^*\mathcal{F})$ if and only if f factors through X_0 . Consider $\mathcal{G} := \operatorname{coker}(\varphi)$ and the exact sequence of $\mathcal{O}_X$ -modules

$$(3) \quad \mathcal{F} \xrightarrow{\varphi} \mathcal{F} \rightarrow \mathcal{G} \rightarrow 0.$$

We claim that $f^*(\varphi)$ is an automorphism if and only if $f^*(\mathcal{G}) = 0$: as $f^*((3))$ is exact, necessity is obvious. If, conversely, $f^*(\mathcal{G}) = 0$ then for any $y \in Y$ $f^*(\varphi)_y$ is a surjective endomorphism of the finitely generated $\mathcal{O}_{Y,y}$ -module $\mathcal{F}_y$, hence is an isomorphism, hence so is $f^*(\varphi)$.

So the sought for $X_0 \subseteq X$ is the complement of the support of $\mathcal{G}$ which is open, because $\mathcal{G}$ is finitely presented.

Proof of lemma 4: Given a scheme X and a short exact sequence $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ of quasi-coherent $\mathcal{O}_X$ -modules with $\mathcal{F}'$ finitely presented and $\mathcal{F}''$ locally free and some $\alpha \in \operatorname{Aut}_{\mathcal{O}_X}(\mathcal{F})$, we have to show the representability by a closed sub-scheme of X of the following functor on X -schemes:

$$F(Y \xrightarrow{f} X) := \begin{cases} * & , \quad f^*(\alpha)(f^*\mathcal{F}') \subseteq f^*\mathcal{F}' \\ \emptyset & , \quad \text{otherwise} . \end{cases}$$

Clearly, F is a Zariski sheaf, so the problem is local on X , i.e. we can assume that $X = \operatorname{Spec}(A)$ is affine, $\mathcal{F}''$ corresponds to some A^n , $\mathcal{F}'$ corresponds to some A -module M for which there is a presentation $A^a \rightarrow A^b \rightarrow M \rightarrow 0$ and $\mathcal{F}$ corresponds to some A -module N . The exact sequence $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ then becomes an exact sequence $0 \rightarrow M \xrightarrow{\iota} N \xrightarrow{\pi} A^n \rightarrow 0$ of A -modules and we are given some $\alpha \in \operatorname{Aut}_A(N)$. Consider $\nu := \pi\alpha\iota$. As all the above sequences are exact after *any* base-change, we have $F(Y \xrightarrow{f} X) \neq \emptyset \iff f^*(\nu) = 0$.

We have a diagram (defining ψ):

$$\begin{array}{ccccccc} A^a & \longrightarrow & A^b & \longrightarrow & M & \longrightarrow & 0 \\ & & & \searrow \psi & \downarrow \nu & & \\ & & & & A^n & & \end{array}$$

which is exact after any base-change, hence $f^*(\nu) = 0 \iff f^*(\psi) = 0$, for any $f : Y \rightarrow X$. So the closed sub-scheme of X we are looking for is the one defined by the ideal of A generated by the coefficients of any matrix representation of ψ .

Proof of 2') $\Rightarrow$ 1) in theorem 1: Under the notations of 1.1 we assume that $\underline{\operatorname{End}}_{\mathcal{F}}$ is representable by a scheme. Then so is $\underline{\operatorname{Aut}}_{\mathcal{F}}$ by lemma 3, hence $\mathcal{F}$ is locally free by what has been shown in 2.2.

2.4

Proof of 3) $\Rightarrow$ 1) and 3') $\Rightarrow$ 1) in theorem 1: Under the notations of 1.1 we assume that X is locally noetherian and that either 3) or 3') holds as well as, by contradiction, that $\mathcal{F}$ is not locally free. By lemma 3 we know in either case that $\underline{\operatorname{Aut}}_{\mathcal{F}}$ is representable by an algebraic space. Using the last assertion of lemma 1 we can assume $X = \operatorname{Spec}(A)$ with A artin local. Then $\underline{\operatorname{Aut}}_{\mathcal{F}}$ is representable by a scheme according to [K], p. 25, 7) contradicting what we proved in 2.2.

Acknowledgements. I would like to thank H. Frommer and M. Volkov for interesting discussions and G. Weckermann for excellent type-setting.

References

- [BLR] S. Bosch, W. Lütkebohmert, M. Raynaud, Néron Models, Ergebnisse der Mathematik, 3. Folge, Band 21, Springer, Heidelberg 1990.
- [K] D. Knutson, Algebraic Spaces, Springer LNM 203.
- [M] H. Matsumura, Commutative ring theory, Cambridge studies in advanced mathematics 8, 1997.
- [N] N. Nitsure, Representability of Gl_E , arXiv:math.AG/0204047.

Mathematisches Institut der WWU Münster
Einsteinstr. 62
48149 Münster
Germany
e-mail: naumannn@uni-muenster.de